

Metal-Insulator oscillations in a Two-dimensional Electron-Hole system

R.J. Nicholas, K. Takashina, M. Lakrimi, B. Kardynal, S. Khym, N.J. Mason
*Department of Physics, Oxford University,
 Clarendon Laboratory, Parks Rd., Oxford, OX1 3PU, U.K.*

D.M. Symons, D. K. Maude and J.C. Portal
*Grenoble High Magnetic Field Laboratory, Max Planck Institut für
 Festkörperforschung and Centre National de la Recherche Scientifique
 BP166, 38042 Grenoble, Cedex 9, France.
 (February 1, 2008)*

The electrical transport properties of a bipolar InAs/GaSb system have been studied in magnetic field. The resistivity oscillates between insulating and metallic behaviour while the quantum Hall effect shows a digital character oscillating from 0 to 1 conductance quantum e^2/h . The insulating behaviour is attributed to the formation of a total energy gap in the system. A novel looped edge state picture is proposed associated with the appearance of a voltage between Hall probes which is symmetric on magnetic field reversal.

The quantum Hall effect has become well known since its first observation by von Klitzing et al [1] but still raises a number of fundamental questions, including the role of edge states which ensure the possibility of dissipationless conduction [2] [3]. In this letter we examine the quantum Hall effect and magnetotransport properties of a bipolar system of coupled electrons and holes and demonstrate that such a system shows qualitatively different behaviour to that observed for single carriers. Such systems have generated considerable interest due to the possibilities of gap formation by both excitonic [4] [5] and single particle [6] interactions. The electron-hole system introduces the new possibility that the edge states of the electron and hole systems may interact, breaking the normal quantum Hall conditions. The first observation of quantum Hall plateaux in an electron-hole system by Mendez et al [7] found that conventional plateaux were formed at quantum numbers corresponding to the difference in the occupancies of the electron and hole Landau levels. Subsequently Daly et al [8] found that in superlattices with closely matched electron and hole densities the special case of zero Hall resistance could be observed. In this letter we examine the behaviour of insulating states formed in a structure containing one layer each of electrons and holes which interact via interband mixing. If mixing occurs between edge states a total energy gap may appear for the system leading to the insulating behaviour. Several gaps can result from the mixing between different electron and hole Landau levels and as a result the system displays oscillatory metallic and insulating behaviour as a function of magnetic field and the Hall conductivity follows a binary sequence oscillating from 0 - 1 - 0 conductance quanta.

The samples studied consisted of a single layer of InAs sandwiched between thick layers of GaSb. This system exhibits a broken gap lineup with the conduction band

edge of the InAs 150meV below the valence band edge of the GaSb. The samples are grown by Metal Organic Vapour Phase Epitaxy (MOVPE) and have a relatively low level of impurities so that the majority of charge carriers are created by intrinsic charge transfer from the GaSb layers to the InAs layer. Typical structures are grown onto semi-insulating GaAs followed by $2\mu\text{m}$ of GaSb to achieve lattice relaxation. The active layer of InAs is typically 30nm thick, followed by a 90nm GaSb cap. Fitting the low field magnetoresistance and Hall effect to classical two carrier formulae give carrier densities of the electrons and holes of order $6.5 \times 10^{15} \text{ m}^{-2}$ and $4.5 \times 10^{15} \text{ m}^{-2}$ respectively. Evidence from magnetotransport suggests that the structure is asymmetric due to pinning of the Fermi level E_F at the surface so that all of the holes are on one side of the structure giving Fermi surfaces for the electrons and holes of similar magnitudes [9] [10]. Determination of the carrier mobilities is less straightforward due to the minigap caused by the electron-hole interaction. However, by applying a large parallel magnetic field, which is known to decouple the bands, [10] [11] the mobilities for the decoupled electrons and holes are estimated to be ~ 20 and $1 \text{ m}^2\text{V}^{-1}\text{s}^{-1}$.

Magnetotransport measurements were made using A.C. techniques with a dilution refrigerator and 15T superconducting magnet, and a ^3He cryostat and 50T pulsed field magnet system on wide Hall bars of width 0.5 mm. Measurements were made with the magnetic field in both forward ($B+$) and reverse ($B-$) directions so that the resistances can be separated into symmetric (S) and antisymmetric (A) parts with respect to field reversal:

$$\begin{aligned} R_{xx}^S &= (R_{xx}(B+) + R_{xx}(B-))/2 = \rho_{xx} \times L/W \\ R_{xx}^A &= (R_{xx}(B+) - R_{xx}(B-))/2 \\ R_{xy}^S &= (R_{xy}(B+) + R_{xy}(B-))/2 \\ R_{xy}^A &= (R_{xy}(B+) - R_{xy}(B-))/2 = \rho_{xy} \end{aligned}$$




FIG. 1. The magnetoresistivity components ρ_{xx} and ρ_{xy} measured in steady fields (main figure) and pulsed field sets for temperatures from 40mK to 4.2 K.

The symmetric part of R_{xx} was used with the length to width ratio (L/W) of the bar to give the diagonal resistivity ρ_{xx} and the antisymmetric part of R_{xy} is taken to be the Hall resistivity, ρ_{xy} according to the Onsager relations. The antisymmetric part of R_{xx} is generally found to be negligibly small, and the symmetric part of R_{xy} is usually attributed to the (small) admixing of the diagonal resistivity [12].

Fig. 1 shows a typical series of plots of the magnetoresistivity and Hall resistivity at temperatures down to 40mK using steady fields and 500mK in pulsed fields. The traces show a number of striking features characteristic of two carrier structures [13] [14] and are very different from single carrier systems. There is a very large positive magnetoresistance with relatively weak oscillatory features at low field. At higher fields there are strong minima at 8.5T and 22T and large maxima at 7T, 15T and 45T which increase rapidly with falling temperature. The peak at 15T has a resistivity which increases to several thousand times its zero field value at 150mK. By contrast the Hall resistivity shows stronger oscillations at low fields, but has only one feature which can be attributed to the formation of a quantum Hall plateau at 8.5T which only appears




FIG. 2. The conductivity components σ_{xx} and σ_{xy} calculated from the data in Fig. 1. The figures indicate the expected electron (ν_e) and hole (ν_h) Landau level occupancies.

at the lowest temperatures. Minima occur where ρ_{xy} approaches zero at 15T by 150mK and 45T by 500mK.

The origins of this behaviour are clearer when the resistivity components are converted to conductivity [15]. Fig. 2 shows that σ_{xy} has a low temperature zero above 40T and in the region 10 - 15T, and approaches zero around 7T. Between these fields one conductance quantum of e^2/h is observed at the lowest temperatures in the steady field data, and a peak moving towards this value with falling temperature is observed at around 20T in the pulsed field data. Minima occur in σ_{xx} when ρ_{xx} has both minima and strong maxima where $\rho_{xx} \gg \rho_{xy}$.

An important feature of broken gap systems is that the electron Landau levels start at a lower energy than the hole levels, and as a function of magnetic field the two sets of levels must cross. Due to the finite interband mixing between the valence and conduction bands the levels anticross leading to the formation of a miniband gap [10] [11] [16] [17] [18] [19] [20], shown schematically in fig. 3i. The movement of the levels through each other leads to oscillatory carrier densities due to the varying density of states, but the position of the Fermi level varies much less than in single carrier systems due to charge transfer between the bands. For structures with a small net doping (i.e. $n_e - n_h$ is small and constant) the Fermi level can lie close to the centre of either electron or hole Landau levels or within the localised tail states of either or both carrier types. In the simplified schematic fig. 3i the Fermi level is shown as constant. The net occupancy of the levels will oscillate between zero around positions a (where $\nu_e - \nu_h = 0 - 0$) and c (where $\nu_e - \nu_h = 1 - 1$) and 1 for b (where $\nu_e - \nu_h = 1 - 0$) and d (where $\nu_e - \nu_h$




FIG. 3. (i) and (ii) the electron and hole Landau levels as a function of magnetic field and spatial position respectively. Electron-like levels are shown in heavier lines. (iii) and (iv) show possible edge current paths in the insulating state with and without the presence of disorder.

$= 2 - 1$). The particularly unusual features of the conduction process occur when the Fermi level lies between equal numbers of electron and hole Landau levels. In this case the system is able to show a complete energy gap which leads to the appearance of oscillatory insulating behaviour as a function of magnetic field.

In single carrier systems the quantum Hall condition also corresponds to a situation where the Fermi level lies within a mobility gap between Landau levels. As the levels approach the sample edges, however, they move upwards in energy and form non-dissipative conducting edge states which generate zeroes in the resistivity and quantum Hall plateaux. The picture can be quite different in interacting electron-hole systems. Considering a spatial plot of the energy levels of the electrons and holes shown in fig. 3ii we expect that as they approach the edges of the structure the electron levels will move upwards due to the edge confinement, while in contrast the hole levels will move downwards. As a result the electron and hole levels will always approach each other but due to the interband coupling the levels will anticross, producing an energy gap. When the Fermi level in the bulk lies between an equal number of electron and hole levels the structure will always show completely insulating be-




FIG. 4. The experimentally measured Hall resistance components $R_{xy}(B+)$ and $R_{xy}(B-)$, and the values deduced for the symmetric and asymmetric Hall resistance R_{xy}^S and R_{xy}^A measured at 100mK. The inset shows the ratio of the experimentally measured Hall resistivities on reversing magnetic field direction.

haviour. This conclusion is obvious when the Fermi level in the bulk also corresponds to the anticrossing gap at the edges, e.g. for $E_F = E_1$ in Fig 3ii. It also holds when it crosses the interacting electron and hole edge states since the number of these states is equal leading to zero net current flow as for $E_F = E_2$ in Fig 3ii. At contacts to the structure we expect edge-states to be at the potential of the originating contact. Although the electron and hole states represent one dimensional current paths with opposite directions (fig. 3iii), we might expect that the edges would be equipotentials giving zero resistivity. This is not the case however, since fluctuations in the potential could cause the gap to rise above the Fermi level, thus disconnecting the channels into two U-turns. More than one such disconnection will lead to a series of isolated conducting loops along the sample edge. One possible arrangement is shown schematically in fig. 3iv.

By contrast when there is population within the bulk of an unequal number of Landau levels there is population of a finite net number of edge states. This gives metallic behaviour with compensated quantum Hall plateaux in the Hall resistance and zeros in ρ_{xx} . There is a rotation through $\sim 90^\circ$ of the equipotentials which lie along the conducting edge states of the bar in the metallic quantum Hall state but were mainly across the bar for the insulating state where no Hall field was generated.

The oscillations between insulating and metallic behaviour occur as the Fermi level moves through regions b,d (metallic) and a,c (insulating) in fig. 3.i. The appearance of an insulating state is crucially linked to the anti-crossing levels and it is interesting to note that a previous report of a re-entrant insulating phase in p-SiGe was tentatively linked to unusual Landau level degeneracy [12] where states may be crossing.

Evidence of a strong distortion of the cu in the insulating state comes from the large component (about 50% of ρ_{xx}) in R_{xy} occurring at the fields where the system becomes insulating. Fig. 4 shows R_{xy} as measured with variations of the magnetic field. The symmetric and antisymmetric parts are also shown. The inset shows $R_{xy}(B+)/R_{xy}(B-)$. Up to ~ 9 T R_{xy} is completely antisymmetric and the ratio is -1. The Hall component has almost no admixing ($\leq 0.5\%$) of the diagonal resistivity ρ_{xx} . At the onset of the insulating state it switches behaviour to become almost totally symmetric. The antisymmetric Hall resistance tends to a large finite voltage is developed across the sample does not depend on the polarity of the applied field thus giving a large value of R_{xy}^S . This has no obvious functional relation to the diagonal resistivity R_{xx}^S which rules out a simple admixing origin. Potential probes, although physically opposite to an accuracy of within $10\mu\text{m}$, are not equipped with the fact that $R_{xy}^A=0$.

Another striking feature of the symmetric Hall resistance is the reproducible resistance fluctuations which are accurately repeated for $B+$ and $B-$. These are shown in the inset to fig. 5, defined by $\Delta R_{xy}^S = R_{xy}^S - \overline{R}_{xy}^S$ is the smoothed background. The function of the resistance fluctuations remains essentially independent of temperature, only increasing in magnitude contrast the Hall resistivity ($=R_{xy}^A$) shows no fluctuations suggesting that any such disorder induced paths are almost completely symmetric with respect to reversal of the direction of the carrier orbit. The violation of the Onsager relations implies the non-local nature of the symmetric Hall voltage.

The temperature dependence of the resistance components ρ_{xx} , R_{xy}^S and ΔR_{xy}^S are shown in fig.5 for the insulating region. The essentially non-Onsager form of the symmetric R_{xy} component leads us to analyse the directly measured resistance rather than resistivity. The diagonal resistivity varies relatively slowly and saturates below ~ 100 mK. The R_{xy}^S and its fluctuations are more strongly temperature dependent. Fits to these with exponentially activated conduction only work over a factor of about two in temperature range, however simple power laws work from the region of 800 to 200 mK where R_{xy}^S is proportional to T^{-3} and ΔR_{xy}^S to T^{-4} .

In conclusion, we have demonstrated that the electron-hole system oscillates between insulating and conducting states as the magnetic field is increased. A qualitative explanation has been proposed based on the formation of an energy gap due to the anticrossing between the electron and hole edge-states. The insulating states have been shown to have unusual behaviour when the Hall resistance becomes symmetric with respect to field reversal.

Acknowledgements: We are grateful to the UK-




FIG. 5. The temperature dependence of ρ_{xx} , R_{xy}^S and ΔR_{xy}^S , showing power law behaviour for the symmetric components of the Hall resistance. The inset shows ΔR_{xy}^S at 100, 225, 300 and 450 mK, demonstrating that the form of the fluctuations is temperature independent.

EPSRC and the EU for continued support of this work.

-
- [1] K. von Klitzing, G. Dorda and M. Pepper, Phys. Rev. Lett. **45**, 494 (1980).
 - [2] M. Buttiker, Phys. Rev. B. **38**, 9375 (1988)
 - [3] D.B. Chklovskii, B.I. Shklovskii, L.I. Glazman, Phys. Rev. B. **46** 4026 (1992)
 - [4] Y. Naveh and B. Laikhtman, Phys. Rev. Lett. **77**, 900 (1996)
 - [5] P.B. Littlewood and X. Zhu, Phys. Scr. **T68**, 56 (1996)
 - [6] S. de-Leon, L. D. Shvartsman, and B. Laikhtman, Phys. Rev. B **60**, 1861 (1999)
 - [7] E.E. Mendez et al, Phys. Rev. Lett. **55**, 2216 (1985)
 - [8] M.S. Daly et al, Phys. Rev. B **53**, R10524 (1996).
 - [9] D. M. Symons et al, Phys. Rev. B **58**, 7292 (1998).
 - [10] M. Lakrimi et al, Phys. Rev. Lett. **152**, 3034 (1997).
 - [11] M. J. Yang, C.H. Yang, B.R. Bennett and B.V. Shanabrook, Phys. Rev. Lett. **78**, 4613 (1997).
 - [12] R.B. Dunford et al, J. Phys.: Condens. Matter. **9**, 1565 (1997)
 - [13] R.J. Nicholas et al, Physica B **201**, 271 (1994).
 - [14] K.S.H. Dalton et al, Surf. Sci. **156**, 305 (1994).
 - [15] R.J. Nicholas et al, Physica E, in press (1999)
 - [16] M. Altarelli, Phys. Rev. B. **28**, 842 (1983).
 - [17] Jin-Chen Chiang et al, Phys. Rev. Lett. **77**, 2053 (1997).
 - [18] L.J. Cooper et al, Phys. Rev. B **57**, 11915 (1998)
 - [19] S.V. Vasil'ev et al, JEPS Lett. **69**, 343 (1999)
 - [20] C. Petchsingh et al, Physica E, (in press) (1999)